

Denitrification and Nitrous Oxide Emissions in Annual Croplands, Perennial Grass Buffers, and Restored Perennial Grasslands

Javed Iqbal*

Dep. of Agronomy
Iowa State Univ.
Ames, IA 50014

Timothy B. Parkin

USDA-ARS
National Lab. for Agric. and the Environ.
Ames, IA 50014

Matthew J. Helmers

Agricultural and Biosystems Engineering
Iowa State Univ.
Ames, IA 50014

Xiaobo Zhou

Monsanto Company
800 N. Lindbergh Blvd.
St. Louis, MO 63167

Michael J. Castellano

Dep. of Agronomy
Iowa State Univ.
Ames, IA 50014

Inclusion of perennial vegetation filter strips (PFSs) in the toeslope of annual cropland watersheds can decrease NO_3^- -N losses to ground and surface waters. Although PFSs are similar to riparian buffers, the processes responsible for NO_3^- -N removal from PFSs are not well understood. Our objectives were to (i) determine the importance of denitrification as a sink for NO_3^- -N loss from PFSs and (ii) evaluate how PFSs alter the biophysical processes that affect the relative importance of N_2O and N_2 emissions. To address our objectives, we used a coupled field laboratory approach with experimental watersheds that included the following treatments: (i) PFSs covering the bottom 10% of the watershed and an annual corn–soybean crop rotation covering the remaining upslope 90% (PFS); (ii) 100% corn–soybean rotation (CORN); and (iii) 19-yr-old 100% restored native grassland (RNG). In situ N_2O flux rates and laboratory $\text{N}_2\text{O}/(\text{N}_2 + \text{N}_2\text{O})$ ratios were highest in CORN watersheds followed by PFS and RNG watersheds. In contrast, potentially mineralizable C and denitrification enzyme activity (DEA) were highest in PFS and RNG watersheds and lowest in CORN watersheds. Furthermore, there was a negative correlation between $\text{N}_2\text{O}/(\text{N}_2 + \text{N}_2\text{O})$ ratio and DEA. In the laboratory, N_2 fluxes were highest in PFS followed by RNG and CORN. These results indicate that PFS watersheds support greater total denitrification while emitting less N_2O than croplands. Greater potentially mineralizable C in PFS and RNG suggest C availability is an important factor affecting more complete denitrification. These results suggest PFSs function similar to riparian buffers and have potential to reduce NO_3^- -N losses from annual croplands by denitrification to N_2 .

Abbreviations: CORN, corn–soybean rotation; DEA, denitrification enzyme activity; DNRA, dissimilatory nitrate reduction to ammonium; NSNWR, Neal Smith National Wildlife Refuge; PFSs, perennial vegetation filter strips; PVC, polyvinyl chloride; RNG, restored native grassland; SOM, soil organic matter; TN, total N; TOC, total soil organic C; VWC, volumetric water content; WFPS, water-filled pore space.

Grassland to cropland conversion has increased N losses to air and water resources. Croplands are a major source of NO_3^- -N flux to rivers and estuaries (Howarth et al., 2012), and agricultural soil management is the leading anthropogenic source of N_2O to the atmosphere (Smith et al., 2007). Many strategies have been developed to mitigate these N losses. Some prominent options include drainage water management (Gilliam et al., 1979), catch (cover) crops (Thorup-Kristensen et al., 2003), and perennial vegetation riparian buffers (Lowrance et al., 1984; Peterjohn and Correll, 1984; Hill, 1996; Hefting et al., 2005).

Supplemental material is available online for this article.

Soil Sci. Soc. Am. J.
doi:10.2136/sssaj2014.05.0221
Received 30 May 2014.

*Corresponding author (jqbal.soil@yahoo.com).

© Soil Science Society of America, 5585 Guilford Rd., Madison WI 53711 USA

All rights reserved. No part of this periodical may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher. Permission for printing and for reprinting the material contained herein has been obtained by the publisher.

A recent variation of perennial vegetation riparian buffers is small-scale incorporation of PFSs in the toeslope portion of cultivated watersheds where the water table approaches the soil surface. This practice has effectively reduced soil erosion and dissolved N losses from upslope croplands (Helmert et al., 2005). For example, Zhou et al. (2010) found a significant decrease in NO_3^- -N concentrations in the vadose zone and groundwater after incorporation of PFSs in corn–soybean cropping systems that are the subject of this study. Others have found that PFSs reduce NO_3^- -N concentrations in surface runoff (Lin et al., 2007; Ryder and Fares, 2008).

Reduced NO_3^- -N concentrations after PFS incorporation into row crop systems can result from a number of processes including microbial N immobilization in soil organic matter (SOM), dissimilatory nitrate reduction to ammonium (DNRA), plant uptake, or denitrification (Matheson et al., 2002; Hefting et al., 2005). Microbial NO_3^- -N immobilization and transfer to SOM is unlikely to be a significant NO_3^- -N sink in cropland riparian buffers (Groffman et al., 1996). In the watersheds that are the subject of this study, less than 10% of isotopically labeled spring-applied $^{15}\text{NO}_3^-$ -N was recovered in SOM in the subsequent fall (Mitchell et al., 2014). Although DNRA may be an important NO_3^- -N sink (Fazzolari et al., 1998), DNRA represents a very short-term sink because the NH_4^+ product is subject to nitrification. Plant uptake can be a significant NO_3^- -N sink in the riparian zone (Peterjohn and Correll, 1984; Hefting et al., 2005; Harrison et al., 2012), but plants represent a relatively finite N pool that can be rapidly remineralized after senescence (Reddy and Patrick, 1984).

In contrast to these relatively small or short-term NO_3^- -N sinks, denitrification is thought to be the major process responsible for NO_3^- -N removal when plant uptake is low (Haycock and Pinay, 1993). However, denitrification can produce several N gas products including the largely inert gas N_2 as well as the greenhouse gas N_2O . Thus, there is significant concern that riparian buffers and PFSs may increase N_2O losses while decreasing NO_3^- -N losses, thereby trading a water quality problem for a greenhouse gas problem (Prather et al., 1995). Indeed, in a riparian zone, Hefting et al. (2006) found increased N_2O emissions in locations with high NO_3^- -N removal efficiency.

Nevertheless, there are number of factors that affect denitrification rates and N gas product ratios (i.e., $\text{N}_2\text{O}/\text{N}_2$). Water, through its effect on O_2 diffusion into the soil and N gas diffu-

sion out of the soil, is well known to affect denitrification rate and $\text{N}_2\text{O}/\text{N}_2$ product ratios. As gas diffusion and O_2 availability decreases, denitrification rate increases and the $\text{N}_2\text{O}/\text{N}_2$ product ratio decreases (Davidson et al., 2000). The availability of potentially mineralizable C and NO_3^- -N can also affect denitrification rate and $\text{N}_2\text{O}/\text{N}_2$ product ratio (Firestone and Davidson, 1989). Potentially mineralizable C can promote denitrification by increasing O_2 consumption (anaerobicity) and substrate availability (when C is limiting). Moreover, high potentially mineralizable C/ NO_3^- -N ratios favor a low $\text{N}_2\text{O}/\text{N}_2$ denitrification product ratio (Firestone and Davidson, 1989). Consistent with these concepts, N_2O fluxes typically decrease with increasing soil C/N ratio (Klemetsson et al., 2005).

Relative to croplands, riparian buffers, and PFSs can alter these biophysical controls on denitrification in many ways. Perennial grasses can affect the water balance by increasing infiltration, soil water holding capacity, and evapotranspiration (Hickman et al., 2010). Perennial grasses can also affect denitrification substrate availability (by both reducing NO_3^- -N and increasing C) and O_2 availability by increasing potentially mineralizable carbon (Pérez-Suárez et al., 2014). Given the many ways in which PFSs can alter biophysical controls on denitrification, as well as the relatively recent implementation of PFSs compared with riparian buffers, there is a critical need for the evaluation of mechanisms responsible for NO_3^- -N removal and N gas production from PFSs.

Using a paired watershed study in Iowa, we investigated the effects of PFSs on NO_3^- -N removal, N_2O emissions, and denitrification. Our study included (i) three watersheds with PFSs covering the bottom 10% of the watershed and corn covering the remaining 90% of the watershed (PFS), (ii) three watersheds with 100% corn (CORN), and (iii) two watersheds with 19-yr-old 100% RNG (Fig. 1). Using comparisons of measurements in perennial filter strip portions of PFS and similar locations in CORN and RNG watersheds, our main objectives were to (i) determine if PFSs are a larger source of N_2O than CORN, (ii) determine if potential denitrification is greater in PFS than CORN, and (iii) quantify the ability of PFS to support complete denitrification of NO_3^- -N to N_2 . Preliminary data demonstrated that PFS installation significantly reduced vadose zone and groundwater NO_3^- -N concentrations (Zhou et al., 2010). Accordingly, we hypothesized that total denitrification is greater, but the $\text{N}_2\text{O}/\text{N}_2$ ratio of denitrification gas products is lower in PFS and RNG watersheds than CORN.

MATERIALS AND METHODS

Site Description

Watersheds were located at the Neal Smith National Wildlife Refuge in Jasper County, Iowa (NSNWR) ($41^\circ 33' \text{N}$, $93^\circ 16' \text{W}$). The refuge includes annual crop production and restored native grassland. The NSNWR is located on the southern Iowa drift plain that contains steep rolling hills of Wisconsin-age loess on pre-Illinoian glacial till (Prior, 1991). Average annual precipitation is 850 mm, most of which occurs in rain events from May to July. Most soils are highly




Fig. 1. Conceptual design of watersheds.

erodible and classified as Ladoga (Mollic Hapludalf) or Otley (Oxyaquic Argiudolls) soil series with 5 to 14% slopes (Nestrud and Worster, 1979; Soil Survey Staff, 2003).

Experimental Design

Eight small zero order (intermittent in hydrological outflow) watersheds in an unbalanced incomplete block design were distributed across three blocks. Blocks are termed Basswood, Interim, and Orbweaver. The size of the watersheds varied from 0.5 to 3.2 ha, with average slopes ranging from 6.1 to 10.5% (Table 1). Interim and Orbweaver blocks each contained three watersheds with the following treatments: 100% annual crop rotation of corn and soybeans (CORN); 10% perennial filter strip at the toeslope position with 90% annual crop rotation of corn and soybeans upslope (PFS); and 100% perennial restored native grassland vegetation (RNG). Basswood contained only CORN and PFS treatments (Table 1). The lack of an RNG watershed in the Basswood block reduced our statistical power, but the RNGs, provided a valuable reference condition that simulated a native prairie ecosystem of the Midwestern United States. Although the PFS always covered the lowest 10% of the watershed (by area), the size and width of the PFS varied from 27 to 41 m based on the size and shape of the watershed. In this study, only the presence of PFS at each watershed position was considered as a treatment, while the area of PFS was not considered as a treatment because Zhou et al. (2010) did not find a significant effect of PFS area on soil or water quality. Details of watershed treatment establishment were reported by Zhou et al. (2010). In the year of this study (2012), corn was the crop in rotation and anhydrous ammonia was knifed into the row crop areas at a rate of 157 kg N ha⁻¹ on 27 Mar. 2012. Grain yield was measured with a combine monitor and converted to aboveground standing biomass at senescence with the standard 0.5 harvest index. Details of the conversion method are provided by Brown et al. (2014). In all other years when corn was the crop in the rotation, the same N fertilizer rate was applied. No fertilizer is added to the 10% of the PFS watersheds maintained in perennial vegetation. In PFS perennial vegetation strips, aboveground standing biomass at senescence was measured by clipping in three randomly placed 1 m² quadrats. Destructive aboveground biomass clipping was not permitted in the RNG. The RNG was managed with annual spring burning, which is typical for native grassland management in the region. Although the 10% PFS establishment was similar to the RNG, it was mowed and harvested rather than burned because of management constraints.

Pretreatment Data

Before the spring of 2007, PFS and CORN watersheds were maintained in bromegrass (*Bromus L.*) for at least 10 yr without fertilizer application. In the spring of 2007, all watersheds were tilled, and PFS and CORN treatments were installed. From 2005 to 2010, 0 to 15 cm of total soil N and organic C decreased in the bottom 10% of the CORN watersheds by approximately 20% (Pérez-Suárez et al., 2014). Pretreatment soil and vegetation data

Table 1. Watershed description and experimental design.

Watershed	Size, ha	Slope, %	Treatment
Basswood-1	0.53	7.5	10% perennial filter strip
Basswood-6	0.84	10.5	100% annual crops
Interim-2	3.19	6.1	10% perennial filter strip
Interim-3	0.73	9.3	100% annual crops
Interim†	–	–	100% restored native grassland
Orbweaver-1	1.18	10.3	10% perennial filter strip
Orbweaver-3	1.24	6.6	100% annual crops
Prairie†	–	–	100% restored native grassland

† Watershed areas of 100% restored native grasslands are unknown, but fall within the range of other watersheds included in this study.

are not available for the PFS and RNG watersheds. However, data from similar watersheds at the research site demonstrate PFS increased total soil N and organic C by increasing net primary productivity relative to annual crops, retaining eroding soil from upslope locations, and minimizing soil erosion from within the perennial vegetation strips (Helmets et al., 2012; Pérez-Suárez et al., 2014). Data from these watersheds suggest aboveground productivity is similar between PFS perennial vegetation strips and RNG, but C4 grasses rather than C3 grasses are more dominant in RNG watersheds (Pérez-Suárez et al., 2014).

In 2005, lysimeters and groundwater wells were installed at “upslope” and toeslope” landscape positions in watersheds that received PFS and CORN treatments in 2007 (Fig. 1). In PFS watersheds, the upslope wells and lysimeters were located where the interface of crops and perennial vegetation would occur after the onset of cropping in 2006. In CORN watersheds, the bottom 10% of the watershed was identified with a procedure identical to that used for identifying the bottom 10% of PFS watersheds, and the upslope wells and lysimeters were located where the interface between crops and perennial vegetation would occur if a 10% PFS were to be inserted in the CORN watersheds. Toeslope wells and lysimeters were located at the bottommost position of the watershed. To better represent the bottom 10% of each watershed in our measurements, two additional slope positions, sideslope and downslope between the upslope and toeslope, were identified. Note that hydrological flow in these watersheds is intermittent, and no stream channel exists because of cultivation operations in the CORN watersheds and growth of perennial vegetation in the PFS and RNG watersheds. Comparing pre- and post-treatment data from lysimeters and wells, Zhou et al. (2010) reported that vadose zone and groundwater NO₃⁻-N concentrations remained low in PFS watersheds (<2.0 mg NO₃⁻-N L⁻¹) after conversion from bromegrass, but conversion from bromegrass significantly increased vadose zone and groundwater NO₃⁻-N concentrations in CORN watersheds (>10 mg NO₃⁻-N L⁻¹).

Zhou et al. (2010) demonstrated lateral groundwater flow through the bottom 10% of watersheds based on higher water table depth above mean sea level in upslope compared with toeslope wells in all watersheds from 2005 to 2008. During our study period (March–October 2012), the groundwater table depth was measured monthly using a submersible level transmitter (Keller America, Newport News, VA). Groundwater samples were also

drawn each month from each well. Water samples were analyzed for NO_3^- -N concentrations on a Quickchem 2000 Automated Ion Analyzer flow injection system with a 0.2 mg L^{-1} detection limit (Lachat Instruments, Milwaukee, WI).

In Situ Nitrous Oxide Gas Flux Measurements

Nitrous oxide flux measurements were performed between 9:00 and 13:00 h from 27 Mar. to 15 Oct. 2012 (the portion of the year when average daily soil temperature was $>10^\circ\text{C}$). Gas flux measurements were performed approximately every other week with greater sampling frequency in the spring and after precipitation events. Our measurement frequency was designed to determine if there was a land use treatment effect on the mean rate of N_2O emissions, not to determine the cumulative amount of N_2O emissions from each land use. A vented, static closed chamber was used to measure gas flux. The chamber consisted of two parts: a polyvinyl chloride (PVC) ring (inner diameter of 25.2 cm and a height of 9.1 cm.) served as a base and was covered with a PVC lid (inner diameter of 25.2 cm and a height of 11.2 cm.). To reduce the effect of radiative heating during gas sampling, thermal insulation tape was added to the outside of PVC chambers. Four gas flux measurement positions on the landscape were selected (upslope, sideslope, downslope and toeslope; Fig. 1). One week before the start of gas flux measurements, one chamber was installed at each location in PFS and RNG watersheds while two chambers were installed in CORN watersheds to account for the systematic heterogeneity (crop row and inter-row). Although one chamber per landscape location (no sub-replication) limited our ability to identify a landscape effect, it standardized the location of chambers in proportion to the watershed area. The chambers were installed to a depth of 5 cm and only moved at the time of fertilizer application and crop planting. During gas measurements, the PVC base was sealed to the PVC chamber over a period of 30 min. Four gas samples drawn at 0, 10, 20, and 30 min with a 10-mL polypropylene syringe were injected into evacuated glass serum vials and brought to the laboratory for gas analysis within 72 h.

Nitrous oxide was analyzed on a gas chromatograph (Agilent 7890), which was operated with an electron capture detector at 350°C for N_2O detection. Gas species separation was accomplished with stainless steel columns packed with Porapak Q, 80/100 mesh and maintained at 85°C . Carrier gas was 10% CH_4 and 90% Ar. Gas emission rates were calculated from the increase in N_2O concentration over time and these relationships were best fit with a linear model. Soil temperature and volumetric water content (VWC) were measured at 5-cm depth near each chamber on each sampling date. Soil temperature was measured with a digital thermometer ($\pm 0.5^\circ\text{C}$), and soil water content was measured via the dielectric constant with a Dynamax SM150 (Houston, TX) soil moisture sensor ($\pm 3\%$ VWC). Volumetric water content data were converted to water-filled pore space (WFPS) using bulk density data from each site.

Soil Physical and Chemical Analysis

To determine soil NO_3^- -N and NH_4^+ -N concentrations during gas flux measurements, two soil sample cores (2-cm diameter to 10-cm depth) were collected within 25 cm of each chamber. Soil samples were extracted in 2 M KCl (5:1 solution: soil ratio). Extracts were filtered through preleached Whatman 1 filter paper and frozen until analysis. Extract ($\text{NO}_3^- + \text{NO}_2^-$)-N (hereafter NO_3^- -N) and NH_4^+ -N concentrations were measured using the Griess-Ilosvay reaction with vanadium(III) chloride as a reducing agent and the Berthelot reaction, respectively (Hood-Nowotny et al., 2010). Soil pH was measured in 1:5 soil/water solution of fresh soil samples. Particle size was determined with the micropipette method (Miller and Miller, 1987). Total C and N were determined by dry combustion elemental analysis.

Denitrification Enzyme Activity

Denitrification enzyme activity was measured from soils collected at each gas flux location (Fig. 1) at five depth increments: 0 to 20, 20 to 40, 40 to 60, 60 to 80, and 80 to 100 cm in September 2012. Soils were brought to the laboratory and kept at 4°C for 3 d until performance of the DEA assay as described by Smith and Tiedje (1979). During the DEA assay, gas samples were taken at 0, 30, 60, 90, and 120 min, stored in evacuated glass vials, and analyzed for N_2O using gas chromatography as described above.

Two phases with an increase in N_2O concentration in headspace over time were found. Rates obtained from soils incubated during Phase I reflected DEA of the existing bacterial enzymes rather than Phase II, which reflects the period of bacterial growth (Smith and Tiedje, 1979). Denitrification rates were calculated as the rate of N_2O accumulation in the headspace in Phase I (between 0 and 60 min) of incubation.

Denitrification Measurements

We measured denitrification using an intact core incubation with (i) ^{15}N isotopic labeling (Mosier and Klemmedtsson, 1994) and (ii) acetylene inhibition methods (Ryden et al., 1987). We observed an exponential decrease in DEA with depth (results) so these experiments were limited to surface soils (0–15 cm). Soils were collected in 20- by 4.1-cm butyrate soil cores from each in situ gas sampling location (one soil core per chamber). Power analyses based on in situ N_2O flux data indicated that one core per location would be sufficient to detect a treatment effect, while one core per landscape location provided little power to detect a landscape effect. After collection, the butyrate sleeves were fitted with rubber stoppers at each end. The samples were kept in dark at 4°C for 3 d until processed. Before denitrification measurements, cores were saturated overnight by wrapping cheese cloth around base of core and putting the base of the cores in 2 cm of deionized water. The cores and tub of water were maintained at 4°C to minimize microbial activity during the wetting phase. Subsequently, the cores were allowed to come to room temperature ($\sim 24^\circ\text{C}$) and drain to field capacity. Cores were then weighed so that the water content of cores could be

kept constant through all denitrification measurements. The internal gas volume of each core was measured using a pressure transducer (Parkin et al., 1984).

Three denitrification measurements were consecutively made on each soil core. In the first set of denitrification measurements (here afterward as No C₂H₂), the ¹⁵N gas flux technique was accomplished by adding 2 mL of enriched ¹⁵NO₃⁻-N (0.5 mg per mL or 1 mg ¹⁵NO₃⁻-N per core) solution in each soil core through 20-cm side port needles. The solution was injected throughout each core to achieve uniform ¹⁵N labeling. Water-filled pore space adjustments were not considered because the soil cores were assumed at field capacity as stated above. Labeling for ¹⁵NO₃⁻-N resulted in 9 to 96% atom percent enrichment in cores with lower values in nitrate-rich soils sampled from the CORN watersheds and higher values in relatively nitrate-poor soils sampled from the PFS and RNG watersheds. The standardized 1-mg ¹⁵NO₃⁻-N addition to each core was chosen based on nearby soil NO₃⁻-N concentrations at the time of soil core sampling and resulted in final NO₃⁻-N concentrations that were within the observed range during field measurements of N₂O flux. After NO₃⁻-N addition, CORN treatment concentrations ranged from 8.7 to 49.5 mg NO₃⁻-N kg soil⁻¹, while field observations during N₂O measurements ranged from 0.11 to 300 mg NO₃⁻-N kg soil⁻¹ with a mean concentration of 22.9 mg NO₃⁻-N kg soil⁻¹; RNG treatment concentrations ranged from 4.7 to 8.2 mg NO₃⁻-N kg soil⁻¹, while field observations during N₂O measurements ranged from 0.03 to 15.2 mg NO₃⁻-N kg soil⁻¹ with a mean concentration of 1.7 mg NO₃⁻-N kg soil⁻¹; PFS treatment concentrations ranged from 5.1 to 40 mg NO₃⁻-N kg soil⁻¹, while field observations during N₂O measurements ranged from 0.05 to 96 mg NO₃⁻-N kg soil⁻¹ with a mean concentration of 8.6 mg NO₃⁻-N kg soil⁻¹.

After enrichment, the cores were sealed with rubber stoppers and incubated for 45.5 h. Gas samples were drawn from the headspace at 0, 4, 21.5, and 45.5 h to ensure a linear increase of N₂O with time; after 0, 4, and 21.5 h samples, 20 mL of laboratory air was added to each core to replace the gas sample. Samples were injected into separate evacuated 10-mL glass serum vials for N₂O and ¹⁵N₂ analyses. After this set of measurements, cores were left open to remove accumulated N₂O, and stoppers were changed for the next set of measurements.

In the second set of measurements (hereafter “Low C₂H₂”), 11 mL of 65 Pa C₂H₂ concentration was added to each soil core with gas volume of 119 mL to yield a final C₂H₂ concentration of approximately 6 Pa which is known to inhibit nitrifier activity and thus N₂O production from nitrification (Mosier, 1980). Cores were then sealed with stoppers and incubated for 29 h. Gas samples were drawn from the headspace at four sampling points (0, 5, 22, and 29 h). Before collecting gas samples at each sampling point, 11 mL of low C₂H₂ was added to each core to replace the withdrawn sample and return the C₂H₂ concentration to 6 Pa. The added air in core was mixed by repeated pumping with a 60-mL syringe, and then 11 mL of gas samples were taken for

N₂O in 6-mL evacuated glass serum vials. After this set of measurements, cores were left open to remove accumulated N₂O, and stoppers were changed for the next set of measurements.

In third set of measurements (hereafter “High C₂H₂”), 11 mL of 100% C₂H₂ gas was added in each soil core of gas volume 119 mL to achieve final concentration of approximately 10 kPa to inhibit the activity of nitrous oxide reductase (Ryden et al., 1987). After sealing with stoppers, the cores were incubated for 28 h. Gas samples were drawn from the head space at four sampling points (0, 5, 23, and 28 h). Before collecting gas samples, 11 mL of C₂H₂ was added to each core, mixed by repeated pumping with a 60-mL syringe, and then 11 mL of gas samples were drawn for N₂O in 6-mL evacuated glass serum vials. The three denitrification measurements were completed in 9 d, and subsequently the soil was air dried and analyzed for residual total ¹⁵N by using a Fisons NA 1500 Series 2 Elemental Analyzer coupled with a Finnigan Delta V mass spectrometer (Thermo Electron, Bremen, Germany). Although C availability is expected to decrease during the incubation, the high water content of soil cores was expected to minimize heterotrophic respiration. By repeating measurements on the same cores, we minimized spatial variability but increased temporal variability. During the three sets of denitrification measurements, N₂O gas production rates were calculated from the increase in N₂O gas concentration from the initial three points, while the last sampling point was discarded.

We measured ¹⁵N₂ and ¹⁵N₂O by using a Finnigan GasBench II (Thermo Electron) interfaced to the mass spectrometer. The analytical precision is 0.2 delta for N₂ and N₂O measurements. Gas chromatography was used to measure total N₂O. In acetylene inhibition methods, N₂O production was measured with linear model based on the increase in concentration over time (accounting for the dilution of laboratory air that was injected to maintain headspace pressure). The following equations from Mosier and Klemmedtsson (1994) were used to calculate N₂ flux from ¹⁵N₂ measurements using the first (time 0) and third sampling point (time 21.5 h):

$$\Delta r = ({}^{29}\text{N}_2/{}^{28}\text{N}_2)_{\text{sample}} - ({}^{29}\text{N}_2/{}^{28}\text{N}_2)_{\text{reference}} \quad [1]$$

$$\Delta t = ({}^{30}\text{N}_2/{}^{28}\text{N}_2 + {}^{29}\text{N}_2)_{\text{sample}} - ({}^{30}\text{N}_2/{}^{28}\text{N}_2 + {}^{29}\text{N}_2)_{\text{reference}} \quad [2]$$

$${}^{15}\text{X}_{\text{N}} = 2.015 (\Delta t / \Delta r) / \{1 + [2.015 (\Delta t / \Delta r)]\} \quad [3]$$

$$d = \Delta t / ({}^{15}\text{X}_{\text{N}})_2 \quad [4]$$

$$\text{Total denitrification N gas evolved from the soil into the core} \\ = \text{total N}_2 \text{ in the core volume} \times d \quad [5]$$

$$\text{N}_2 \text{ flux} = \Delta C / A t \quad [6]$$

where “sample” is an air sample collected within the core at a time *t* after addition of ¹⁵NO₃⁻-N, “reference” is an ambient laboratory air sample at the time of the experiment, (²⁹N₂/²⁸N₂) and (³⁰N₂/²⁸N₂ + ²⁹N₂) are ion current ratios determined by the mass spectrometer, ¹⁵X_N is mole fraction of ¹⁵N in the soil NO₃⁻ pool, *d* is the fraction of total N gas in the soil core attrib-

Table 2. Mean (\pm standard error) groundwater depth and NO_3^- -N concentration.†

	CORN			PFS		
	Groundwater height above mean sea level*	Groundwater NO_3^- -N ^{NS}	Soil NO_3^- -N*	Groundwater height above mean sea level ^{NS}	Groundwater NO_3^- -N*	Soil NO_3^- -N*
	m	mg L ⁻¹	mg kg ⁻¹	m	mg L ⁻¹	mg kg ⁻¹
Upslope	298.9 \pm 4.03	4.63 \pm 0.57	26.94 \pm 4.16	298.4 \pm 2.09	3.78 \pm 1.08	9.75 \pm 1.32
Toeslope	289.4 \pm 4.87	5.05 \pm 1.79	12.54 \pm 1.60	289.4 \pm 6.00	1.58 \pm 0.79	6.13 \pm 1.77

† Data represent mean values of six sampling dates from April to September 2012. Mean (\pm standard error) NO_3^- -N concentration in surface soils (0–10-cm depth) represent mean values of 14 samplings dates; data represent mean values of 14 sampling dates from March to October 2012. Treatments include 100% annual row crop (CORN) and 10% perennial filter strip (PFS) watersheds at Neal Smith National Wildlife Refuge ($N = 3$). The Restored native grassland did not contain wells. See Fig. 1 for sample locations. Probability from paired t test between upslope and toeslope; * $P < 0.01$; ^{NS}, Not significant.

unable to denitrification, A is soil surface area in the soil core, t is the time when core was sealed, and ΔC is the change in concentration of $^{30}\text{N}_2$ and $^{29}\text{N}_2$ in the soil core during time t .

By using data from the acetylene inhibition methods, we calculated N_2 production as the difference in N_2O production between cores with “No C_2H_2 ” and “High C_2H_2 ” (Ryden et al., 1979). Similarly, $\text{N}_2\text{O}/(\text{N}_2 + \text{N}_2\text{O})$ was calculated from estimates of N_2O produced from intact cores with No C_2H_2 and High C_2H_2 . Estimates of N_2O production from denitrification were calculated as N_2O flux with Low C_2H_2 because nitrification was inhibited (Mosier, 1980). We calculated N_2O production from nitrification by taking the difference of N_2O rates in soil cores with No C_2H_2 and Low C_2H_2 . However, values were extremely low, and slightly negative values were common suggesting little to no N_2O production from nitrification, as would be expected at 100% water holding capacity (Zhu et al., 2013).

Carbon Mineralization

Potentially mineralizable C (C_{min}) was determined from soil samples collected during September 2012 for the denitrification measurements. Ten grams of soil was incubated in 120-mL bottles during a 5-d incubation at 25°C. Headspace CO_2 concentration was quantified daily using an LI-7000 infrared gas analyzer (LI-COR, Lincoln, NE). Carbon dioxide emissions were calculated by linear interpolation and numerical integration between sample times.

Statistical Analysis

All data were tested for normality and homogeneity of variance. If needed, data were log transformed before statistical analysis. In CORN watersheds, in situ measurements of N_2O fluxes, NO_3^- -N, and NH_4^+ -N were calculated by taking the average of data from row and inter-row because there were no significant differences between these locations. For in situ field measurements, the effects of treatment, time, landscape, treatment \times time, treatment \times landscape, and treatment \times time \times landscape were evaluated for soil N_2O fluxes, temperature, moisture, and NO_3^- -N and NH_4^+ -N concentration with a repeated measures analysis of variance (ANOVA) model with fixed effects of treatment, time, landscape, and their interactions and block as a random effect (SAS Institute, Cary, NC). We did not calculate cumulative N_2O flux from the 14 in situ measurements because our goal was not to determine the cumulative amount of N_2O emissions from each

land use treatment. For laboratory measurements including DEA, denitrification, C_{min} , and soil properties, the same ANOVA model was used but without repeated measures. Significant main effects were assessed with Tukey’s range test for mean separation. Groundwater height above sea level, groundwater NO_3^- -N, and soil NO_3^- -N mean values from upslope and toeslope locations were compared using paired t tests.

RESULTS

Groundwater and NO_3^- -N Dynamics

Groundwater height above mean sea level decreased from upslope to toeslope wells (Table 2). Mean groundwater NO_3^- -N concentrations in the CORN did not change from upslope and toeslope wells; however, mean groundwater NO_3^- -N concentration in the PFS significantly decreased from the upslope to toeslope wells. A decrease in soil NO_3^- -N (2 M KCl extractions) from upslope to toeslope was observed in both CORN and PFS (Fig. 2 and Table 2).

Soil Properties and Field Measurements

Soil physical properties were not significantly different among treatments. However, total soil organic carbon (TOC), total N, and C/N ratio were significantly higher in PFS and RNG than CORN (Table 3). In CORN and PFS watersheds, average corn aboveground standing biomass at senescence was 14.6 Mg dry matter ha^{-1} (grain + residue). Perennial vegetation aboveground standing biomass at senescence in the perennial filter strips was 6.6 Mg dry matter ha^{-1} .

Across all field measurements, mean soil WFPS did not differ among the treatments (Table 4), but within treatments WFPS increased from upslope to toeslope in PFS and decreased from upslope to toeslope in CORN (Table S1). Mean soil temperature across all measurement dates decreased among treatments in the order of CORN > RNG > PFS (Table 3 and Table 4). Across all measurements, mean soil NH_4^+ -N and NO_3^- -N concentrations were highest in CORN followed by PFS and RNG (Fig. 2).

Across all dates, mean in situ soil N_2O flux rate was highest in CORN but did not differ among PFS and RNG (Fig. 3). On 12 of 14 measurement dates, N_2O flux in CORN was greater than PFS and RNG (Fig. 3). There was a treatment by landscape position interaction on N_2O (Table 4). Within treatments, N_2O fluxes in PFS decreased from upslope to toeslope. In contrast, N_2O fluxes in CORN increased from upslope to toeslope (Fig. 3). There was no effect of landscape position on N_2O fluxes in RNG.

Denitrification Enzyme Activity

Across all treatments, DEA was highest in the surface soil (20-cm depth) and exponentially decreased with increasing soil depth (Fig. 4). As a proportion of total DEA from 0 to 100 cm, 69.6% occurred from 0 to 20 cm, 15.0% from 20 to 40 cm, 6.4% from 40 to 60 cm, 5.8% from 60 to 80 cm, and 3.3% from 80 to 100 cm. Landscape position did not affect DEA (Table 5). Opposite in situ N_2O fluxes, DEA was higher in RNG and PFS than CORN (Fig. 4). There was a positive linear correlation between TOC and the natural logarithm of DEA ($R = +0.48$, $P = 0.04$; Fig. 5) as well as NH_4^+-N and DEA ($R = +0.66$, $P < 0.01$; Fig. 5). There was a negative linear correlation between $NO_3^- - N$ and the natural logarithm of DEA ($R = -0.71$, $P < 0.01$; Fig. 5).

Denitrification: C_2H_2 Technique

Across treatments, there was no consistent pattern between denitrification and landscape position. However, we caution that our analysis of landscape position was limited because of a lack of statistical power. Thus, we focus on the treatment effect and correlations among DEA, $NO_3^- - N$ concentration, potentially mineralizable C, and denitrification.

Total denitrification rates ($N_2 + N_2O$), measured in intact cores, were not different among the treatments (Fig. 6 and Table 5), and soil $NO_3^- - N$ concentrations and total denitrification rates were not correlated. However, across treatments, N_2O production from denitrification did not show the same pattern as total denitrification; N_2 production from denitrification was highest in PFS followed by CORN and RNG (Fig. 6). Moreover, the pattern of N_2O production from denitrification was opposite the pattern of DEA and potentially mineralizable C (Fig. 4 vs. Fig. 6 and Fig. 7).

The $N_2O/(N_2 + N_2O)$ ratio was significantly lower in PFS than RNG and CORN (Fig. 6). This was also opposite the pattern of DEA and potentially mineralizable C. There was a negative correlation between $N_2O/(N_2 + N_2O)$ and the natural logarithm of DEA (Fig. 8) and a positive exponential relationship between $N_2O/(N_2 + N_2O)$ and $NO_3^- - N$ (Fig. 8).

Denitrification: ^{15}N Technique

Total N_2 ($^{15}N_2 + ^{14}N_2$) production was highest in PFS followed by RNG and CORN (Fig. 9). Across treatments, the pattern in total N_2 production was coincident with DEA and potentially mineralizable C, but opposite several N_2O measurements including N_2O fluxes measured in situ, N_2O production during C_2H_2 inhibition, and N_2O production during the ^{15}N tracing (Fig. 3, 6, and 9). Residual ^{15}N remaining in the intact soil cores after incubation exhibited the opposite pattern of $^{15}N_2$ fluxes and was higher in CORN than PFS or RNG (Fig. 10). Denitrification measurements with C_2H_2 method were approximately double those measured with the ^{15}N method.




Fig. 2. In situ soil (a) NH_4^+-N and (c) $NO_3^- - N$ contents as determined at four landscape positions (upslope, sideslope, downslope, and toeslope) at 10-cm depth in each treatment during the growing season (March to September) of 2012. Error bars indicate standard error of the mean ($n = 28-42$). Different letters indicate significant differences among landscape positions within each treatment as determined by Tukey's test. Mean in situ soil NH_4^+-N and $NO_3^- - N$ contents across landscape positions are also shown in Fig. 2b and 2d, respectively, where different letters indicate significant treatment differences determined after a significant main effect of treatment in repeated measures ANOVA. PFS, three watersheds with perennial vegetation filter strips covering the bottom 10% of the watershed and corn covering the remaining 90% of the watershed; CORN, three watersheds with 100% corn; RNG, two watersheds with 19-yr-old 100% restored native grassland.

DISCUSSION

Groundwater and $NO_3^- - N$ Dynamics

Groundwater height above mean sea level decreased from upslope to toeslope wells, providing evidence for lateral water flow through the bottom 10% of CORN and PFS watersheds (Table 2). Moreover, the relatively steep slopes (6–10%) of the study wa-

Table 3. Mean (\pm standard error) of selected soil properties of watersheds from 0- to 15-cm soil depth at Neal Smith National Wildlife Refuge.†

Property	PFS	CORN	RNG
Total organic carbon, %	2.99 \pm 0.17a	2.32 \pm 0.09b	2.78 \pm 0.16a
Total nitrogen, %	0.26 \pm 0.0a	0.22 \pm 0.01b	0.24 \pm 0.01ab
C/N ratio	11.32 \pm 0.37ab	10.53 \pm 0.08b	11.71 \pm 0.28a
Bulk density, g cm ⁻³	1.17 \pm 0.04a	1.22 \pm 0.03a	1.18 \pm 0.02a
pH	6.38 \pm 0.18b	6.12 \pm 0.11b	6.91 \pm 0.08a
Sand, %	8.06 \pm 3.00a	9.89 \pm 3.57a	9.53 \pm 1.84a
Silt, %	25.89 \pm 1.17a	24.21 \pm 1.38a	22.39 \pm 1.31a
Clay, %	66.06 \pm 2.61a	65.90 \pm 2.34a	68.08 \pm 1.78a
Soil temperature, °C	17.00 \pm 0.39b	18.11 \pm 0.40a	17.73 \pm 0.45ab
Water-filled pore space, %	54.34 \pm 1.55a	55.19 \pm 1.27a	53.68 \pm 1.68a

† Same letters within a row indicate insignificant differences ($P > 0.05$). PFS, perennial filter strip; RNG, restored native grassland.

Table 4. Probability values (repeated measured ANOVA) of the impact of treatment, landscape, and treatment × landscape interaction on in situ N₂O fluxes and parameters measured during the growing season (March to September) of 2012 (14 sample dates).

Factor	N ₂ O	Soil temperature	Water-filled pore space	NH ₄ ⁺ -N	NO ₃ ⁻ -N
Treatment	<0.01	<0.01	0.26	<0.01	<0.01
Landscape	0.95	0.33	0.64	0.01	<0.01
Treatment × landscape	0.02	0.44	<0.01	0.02	0.02

tersheds (which are not tilled) likely promote lateral hydrological flow paths (Zhu and Lin, 2009), and lateral flow can promote NO₃⁻-N transport downslope (Castellano et al., 2013). The coincident decrease in groundwater levels and NO₃⁻-N concentrations from upslope to toeslope wells in the PFS suggests PFS create a NO₃⁻-N sink during lateral flow through these systems. Consistent with this interpretation of 2012 data presented herein, Zhou et al. (2010) concluded that lateral flow through PFS led to NO₃⁻-N removal from 2005 to 2008 in the watersheds that are the subject of this study.

In Situ Nitrous Oxide Fluxes

Relative to CORN, consistently low N₂O fluxes in PFS that were similar to RNG (Fig. 3) could reflect not only the lack of direct fertilizer application to the PFS portion of the watershed,




Fig. 3. (a) In situ soil N₂O-N fluxes as determined during the growing season (March to September) of 2012 and (b) mean N₂O-N fluxes at four landscape positions (upslope, sideslope, downslope and toeslope) in each treatment. Error bars indicate standard error of the mean. Different letters indicate significant differences among landscape positions within each treatment as determined by Tukey's test. (c) Mean soil N₂O-N flux rates across landscape positions within treatments. Different letters indicate significant treatment differences determined after a significant main effect of treatment in repeated measures ANOVA.

but also (i) fewer N₂O emissions due to nitrification, (ii) differences in corn vs. perennial vegetation N sinks, and (iii) greater N₂O reduction (i.e., more complete denitrification). Differences in nitrification N₂O emissions could partially explain the differences because CORN watersheds received 157 kg ha⁻¹ of anhydrous ammonia fertilizer. Nevertheless, during the course of field measurements (March–September), mean soil NH₄⁺-N concentration was similar among CORN and PFS despite the large NH₄⁺-N addition to CORN, indicating greater N mineralization in PFS due to higher SOC and TN. These data suggest reduction differences in N₂O fluxes are more likely due to differences in NO₃⁻-N rather than NH₄⁺-N substrate.

Although N inputs to the lower 10% of CORN and PFS watersheds differ, so do vegetation N sinks. Total aboveground dry matter production of 14.6 Mg ha⁻¹ in CORN vs. 6.6 Mg ha⁻¹ in the perennial vegetation portion of PFS suggest that corn is a greater N sink than perennial vegetation. However, the temporal pattern of N uptake differs among these plants so that corn N uptake is concentrated in the summer, whereas prairie vegetation N uptake begins in the early spring (Abendroth et al., 2011; Pérez-Suárez et al., 2014). The lack of spring N demand in CORN watersheds may help to explain why treatment differences in N₂O flux were greatest in the spring and decreased over time (Fig. 3).

Potentially mineralizable C may also be an important factor affecting treatment differences in N₂O flux, particularly in the early spring when vegetation productivity was low in all treatments and CORN N₂O emissions were approximately 100% greater than PFS. Across treatments, patterns of N₂O fluxes and N₂O/(N₂O + N₂) ratios were opposite potentially mineralizable C, suggesting differences in C availability may affect the magnitude of N₂O fluxes and the N₂O/(N₂O + N₂) ratio (Fig. 3 and 6). Low mineralizable C availability can limit the reduction of NO₃⁻-N to N₂O and subsequently N₂O to N₂ (e.g., Parsons et al., 1993; Mitchell et al., 2013). Accordingly, greater




Fig. 4. (a) Denitrification enzyme activity (DEA) measured from 0- to 100-cm depth in three treatments. Analysis of variance results ($P > F$) for treatment, depth, and treatment × depth are shown. (b) Denitrification enzyme activity measured at 0- to 15-cm depth in three treatments. Different letters indicate significant differences among treatments as determined by Tukey's test after a significant main effect of treatment ($P < 0.01$). Vertical error bars indicate standard error of the mean ($n = 8-12$).

Table 5. Probability values (two way ANOVA) of the impact of treatment, landscape, and treatment × landscape interaction on parameters measured in the laboratory.†

Factor	DEA	Cmin	Acetylene inhibition experiment				¹⁵ N tracer experiment		
			N ₂ O‡	N ₂	N ₂ + N ₂ O	N ₂ O/(N ₂ + N ₂ O)	¹⁵ N ₂	N ₂ O§	Residual ¹⁵ N
Treatment	<0.01	<0.01	0.28	0.04	0.35	0.05	0.04	0.10	<0.01
Landscape	0.83	0.20	<0.01	0.73	0.52	0.32	0.87	0.04	0.38
Treatment × landscape	0.96	0.28	0.46	0.78	0.76	0.59	0.65	0.46	0.30

† DEA, denitrification enzyme activity; Cmin, potentially mineralizable carbon.

‡ Denitrified N₂O from the acetylene inhibition experiment.

§ Total N₂O from the ¹⁵N tracer experiment.

organic C availability in PFS could have resulted in greater N₂O reduction to N₂. This result is consistent with the C₂H₂ and ¹⁵N laboratory measurements that indicated similar total denitrification across treatments but lower N₂O production in PFS. In RNG, high C availability and low NO₃⁻-N availability due to the lack of upslope N fertilization and accompanying NO₃⁻-N runoff may have contributed to low N₂O flux.

Denitrification, Denitrification Enzyme Activity, and the N₂O/(N₂ + N₂O) Ratio

The DEA assay evaluates denitrification potential by providing nonlimiting conditions for a short time period. Our results indicated that despite lower NO₃⁻-N concentrations, soils in PFS and RNG had greater denitrification potential than those in CORN. These results are consistent with higher DEA in riparian buffers compared with crop fields (Groffman et al., 1993; Kim, 2008).

Consistent with previous work (Davis et al., 2008), we also observed a positive correlation between TOC and DEA as well as NH₄⁺-N and DEA. The positive relationship between NH₄⁺-N and DEA could be due to (i) NH₄⁺-N regulation of NO₃⁻-N consumption and production (Davis et al., 2008) or (ii) it can be an indirect relationship resulting positive correlations among TOC, NH₄⁺-N, potentially mineralizable C, and N mineralization (Schomberg et al., 2009; Haddad et al., 2013).

Opposite patterns of in situ N₂O fluxes and DEA across the watershed treatments led us to pursue denitrification measurements. Despite higher NO₃⁻-N concentrations and in situ N₂O emissions, less N₂ production was found in CORN than PFS soils. This pattern was consistent with lower DEA, lower potentially mineralizable C, and lower TOC in CORN compared with PFS watersheds. A number of factors control the production and consumption of N₂O and N₂ through the denitrification process. The complete reduction of NO₃⁻-N to N₂ through denitrification is controlled by the relative availability of oxidant (oxidized N) and reductant (most commonly organic C). If oxidant availability greatly exceeds reductant availability, then the oxidant may be incom-

pletely utilized; that is, N₂O rather than N₂ will be produced (Firestone and Davidson, 1989).

Lower NO₃⁻-N concentrations, higher organic C, and higher DEA in PFS might have stimulated the reduction of N₂O to N₂, whereas higher NO₃⁻-N availability and lower C availability in CORN likely limited N₂O reduction (Fig. 2, 3, and 6). Across all treatments, DEA was negatively associated with N₂O/(N₂ + N₂O) while NO₃⁻-N was positively associated with N₂O/(N₂ + N₂O) (Fig. 8). Similar to our results, Hill et al. (2000) observed a significant increase in denitrification after in situ additions of labile C in a riparian zone. With regard to denitrification, these results suggest perennial filter strips function similarly to riparian buffer zones (Lowrance et al., 1984; Matheson et al., 2002) and wetlands (Harrison et al., 2012).

Consistent with previous research, results from the ¹⁵N and C₂H₂ denitrification measurements were well correlated (Myrold, 1990). Although the C₂H₂ method yielded denitrification rates that were approximately double those measured with ¹⁵N (~2.5 vs. 5.0 mg N g⁻¹ soil h⁻¹), these results are reasonably close given inherently high variability in these methods. Denitrification is well known to be highly spatially and temporally variable (Parkin, 1987; Groffman et al., 2009). In our laboratory analyses we could not eliminate both sources of variation because of sample number constraints, so we decided to minimize spatial variation (by conducting the ¹⁵N and C₂H₂ analyses consecutively on identical, intact soil cores). Interestingly, the C₂H₂ analyses yielded higher denitrification rates despite the fact that they were conducted after the ¹⁵N analyses without any substrate addition. This result suggests substrate availability was not limiting denitrification. Although NO₃⁻-N addition can stimulate denitrification and increase the N₂O/(N₂ + N₂O)




Fig. 5. Relationships between the natural logarithm of denitrification enzyme activity (ln DEA) and (a) total organic carbon (TOC mg kg⁻¹), (b) Soil NH₄⁺-N (mg kg⁻¹), and (c) Soil NO₃⁻-N concentrations (mg kg⁻¹).




Fig. 6. Results from the acetylene inhibition experiments (see method details in Methods): (a) Denitrified N_2O , N_2 , and $N_2O + N_2$ and (b) $N_2O / (N_2 + N_2O)$ ratio. Vertical error bars indicate standard error of the mean ($n = 8-12$). Analysis of variance results ($P > F$) for the treatment main effect are shown. Different letters indicate significant differences among treatments as determined by Tukey's test.

ratio, we note that our $^{15}\text{NO}_3^- - \text{N}$ additions were proportionally largest (relative to background $\text{NO}_3^- - \text{N}$) in the RNG treatments which had the lowest $N_2O / (N_2 + N_2O)$ ratio and proportionally smallest in the CORN treatments which had the highest $N_2O / (N_2 + N_2O)$ ratio and lower denitrification rates.

CONCLUSIONS

Our data show that perennial filter strips can decrease $\text{NO}_3^- - \text{N}$ concentrations in near-surface groundwater without increasing N_2O losses to the atmosphere. High potentially min-




Fig. 7. Potentially mineralizable carbon as determined from a 5-d laboratory incubation during September 2012. Vertical error bars indicate standard error of the mean ($n = 8-12$). Analysis of variance result ($P > F$) for the treatment main effect is shown. Different letters indicate significant differences among treatments as determined by Tukey's test.




Fig. 8. Relationships between $N_2O / (N_2 + N_2O)$ ratio and (a) the $\ln \text{DEA}$ and (b) $\text{NO}_3^- - \text{N}$ contents.

eralizable C coupled with lateral flow of $\text{NO}_3^- - \text{N}$ from upslope cropland through downslope PFS likely creates a favorable environment for denitrification. Together, laboratory experiments and field observations suggest greater C availability in PFS than




Fig. 9. Result from (a) the ^{15}N tracer experiment: N_2 fluxes and (b) N_2O fluxes. Vertical error bars indicate standard error of the mean ($n = 8-12$). Analysis of variance results ($P > F$) for treatment main effect are shown. Different letters indicate significant differences among treatments.




Fig. 10. The percentage of ^{15}N remaining in intact soil cores after 9 d of enrichment. Analysis of variance results ($P > F$) for treatment main effect are shown. Different letters indicate significant differences among treatments as determined by Tukey's test. Vertical error bars indicate standard error of the mean ($n = 8\text{--}12$).

CORN leads to more complete denitrification of $\text{NO}_3^- \text{--N}$ to N_2 . Perennial vegetation filter strips can enhance potentially mineralizable C through three processes: (i) PFSs trap sediment that is eroded from upslope organic C-rich surface soils; (ii) PFSs minimize erosion within the PFS buffer; and (iii) perennial grasses produce more belowground net primary productivity than the corn-soybean rotation crop system (Helmers et al., 2012; Pérez-Suárez et al., 2014).

Lower residual ^{15}N in laboratory-incubated cores (Fig. 10) and lower groundwater $\text{NO}_3^- \text{--N}$ concentration (Table 2) in PFS vs. CORN watersheds confirmed that PFSs promote greater $\text{NO}_3^- \text{--N}$ removal efficiency than croplands. Furthermore, relatively low $\text{N}_2\text{O}/(\text{N}_2 + \text{N}_2\text{O})$ ratios in PFS suggest that these systems may have a higher nitrate removal capacity than the load to which they are subjected because reduction in the $\text{N}_2\text{O}/(\text{N}_2 + \text{N}_2\text{O})$ ratio might be expected before an increase in $\text{NO}_3^- \text{--N}$ is transport through PFS because of lower total denitrification. Although the amount of $\text{NO}_3^- \text{--N}$ that can be removed by PFSs is affected by complex interactions among hydrology, vegetation, and soil processes, it appears that the area of PFSs in this study (10% of the watershed) is sufficient to remove large amounts of NO_3^- runoff.

ACKNOWLEDGMENTS

We thank Christopher Witte, Carolina Cordova, and Omar de Kok-Mercado for their assistance in field sampling. Funding for the research site was provided by the Leopold Center for Sustainable Agriculture, Iowa State University College of Agriculture and Life Sciences, USDA Forest Service Northern Research Station, the Iowa Department of Agriculture and Land Stewardship, and the USDA-AFRI Managed Ecosystems program under proposal no. 2011-00512. This research is part of a regional collaborative project supported by USDA-NIFA Award no. 2011-68002-30190, "Cropping Systems Coordinated Agricultural Project (CAP): Climate Change, Mitigation, and Adaptation in Corn-based Cropping Systems." Project web site: sustainablecorn.org.

SUPPLEMENTAL MATERIAL

Supplementary information associated with this article includes in situ water filled pore space (%) and soil temperature ($^{\circ}\text{C}$) as determined across landscape positions in perennial filter strip, 100% annual row crop, and restored native grassland during the growing season of 2012.

REFERENCES

- Abendroth, L.J., R.W. Elmore, M.J. Boyer, and S.K. Marlay. 2011. Corn growth and development. Iowa State Univ., Ames, IA.
- Brown, K.H., E.M. Bach, R.A. Drijber, K.S. Hofmockel, E.S. Jeske, J.E. Sawyer, and M.J. Castellano. 2014. A long-term N fertilizer gradient has little effect on soil organic matter in a high-intensity maize production system. *Glob. Change Biol.* 20:1339–1350. doi:10.1111/gcb.12519
- Castellano, M.J., D.B. Lewis, and J.P. Kaye. 2013. Response of soil nitrogen retention to the interactive effects of soil texture, hydrology, and organic matter. *J. Geophys. Res.* 118:280–290. doi:10.1002/jgrg.20015
- Davidson, E.A., M. Keller, H.E. Erickson, L.V. Verchot, and E. Veldkamp. 2000. Testing a conceptual model of soil emissions of nitrous and nitric oxides. *Bioscience* 50:667–680. doi:10.1641/0006-3568(2000)050[0667:TACMS]2.0.CO;2
- Davis, J.H., S.M. Griffith, W.R. Horwath, J.J. Steiner, and D.D. Myrold. 2008. Denitrification and nitrate consumption in an herbaceous riparian area and perennial ryegrass seed cropping system. *Soil Sci. Soc. Am. J.* 72:1299–1310. doi:10.2136/sssaj2007.0279
- Fazzolari, É., B. Nicolardot, and J.C. Germon. 1998. Simultaneous effects of increasing levels of glucose and oxygen partial pressures on denitrification and dissimilatory reduction to ammonium in repacked soil cores. *Eur. J. Soil Biol.* 34:47–52. doi:10.1016/S1164-5563(99)80006-5
- Firestone, M.K., and E.A. Davidson. 1989. Microbiological basis of NO and N_2O production and consumption in soil. p. 7–21. In: M.O., Andreae, and D.S. Schimel, editors, Exchange of trace gases between terrestrial ecosystems and the atmosphere. John Wiley & Sons, New York.
- Gilliam, J., R. Skaggs, and S. Weed. 1979. Drainage control to diminish nitrate loss from agricultural fields. *J. Environ. Qual.* 8:137–142. doi:10.2134/jeq1979.00472425000800010030x
- Groffman, P.M., K. Butterbach-Bahl, R.W. Fulweiler, A.J. Gold, J.L. Morse, E.K. Stander, C. Tague, C. Tonitto, and P. Vidon. 2009. Challenges to incorporating spatially and temporally explicit phenomena (hotspots and hot moments) in denitrification models. *Biogeochemistry* 93:49–77. doi:10.1007/s10533-008-9277-5
- Groffman, P.M., G. Howard, A. Gold, and W. Nelson. 1996. Microbial nitrate processing in shallow groundwater in a riparian forest. *J. Environ. Qual.* 25:1309–1316. doi:10.2134/jeq1996.00472425002500060020x
- Groffman, P.M., C.W. Rice, and J.M. Tiedje. 1993. Denitrification in a tallgrass prairie landscape. *Ecology* 74:855–862. doi:10.2307/1940811
- Haddad, S.A., M.A. Tabatabai, and T.E. Loynachan. 2013. Biochemical processes controlling soil nitrogen mineralization under waterlogged conditions. *Soil Sci. Soc. Am. J.* 77:809–816. doi:10.2136/sssaj2012.0231
- Harrison, M.D., P.M. Groffman, P.M. Mayer, and S.S. Kaushal. 2012. Nitrate removal in two relict oxbow urban wetlands: A ^{15}N mass-balance approach. *Biogeochemistry* 111:647–660. doi:10.1007/s10533-012-9708-1
- Haycock, N., and G. Pinay. 1993. Groundwater nitrate dynamics in grass and poplar vegetated riparian buffer strips during the winter. *J. Environ. Qual.* 22:273–278. doi:10.2134/jeq1993.00472425002200020007x
- Hefting, M.M., R. Bobbink, and M.P. Janssens. 2006. Spatial variation in denitrification and N_2O emission in relation to nitrate removal efficiency in a N-stressed riparian buffer zone. *Ecosystems* 9:550–563. doi:10.1007/s10021-006-0160-8
- Hefting, M.M., J.-C. Clement, P. Bienkowski, D. Dowrick, C. Guenat, A. Butturini, S. Topa, G. Pinay, and J.T.A. Verhoeven. 2005. The role of vegetation and litter in the nitrogen dynamics of riparian buffer zones in Europe. *Ecol. Eng.* 24:465–482. doi:10.1016/j.ecoleng.2005.01.003
- Helmers, M.J., D. Eisenhauer, M. Dosskey, T. Franti, J. Brothers, and M. McCullough. 2005. Flow pathways and sediment trapping in a field scale vegetative filter. *Trans. ASAE* 48:955–968. doi:10.13031/2013.18508
- Helmers, M.J., X. Zhou, H. Asbjornsen, R. Kolka, M.D. Tomer, and R.M. Cruse. 2012. Sediment removal by prairie filter strips in row-cropped ephemeral watersheds. *J. Environ. Qual.* 41(5):1531–1539. doi:10.2134/jeq2011.0473
- Hickman, G.C., A. Vanloocke, E.G. Dohleman, and C.J. Bernacchi. 2010. A comparison of canopy evapotranspiration for maize and two perennial grasses identified as potential bioenergy crops. *GCB Bioenergy* 2:157–168. doi:10.1111/j.1757-1707.2010.01050.x
- Hill, A.R. 1996. Nitrate removal in stream riparian zones. *J. Environ. Qual.* 25:743–755. doi:10.2134/jeq1996.00472425002500040014x
- Hill, A.R., K. Devito, S. Campagnolo, and K. Sanmugadas. 2000. Subsurface denitrification in a forest riparian zone: Interactions between hydrology

- and supplies of nitrate and organic carbon. *Biogeochemistry* 51:193–223. doi:10.1023/A:1006476514038
- Hood-Nowotny, R., N.H.-N. Umana, E. Inselbacher, P. Oswald-Lachouani, and W. Wanek. 2010. Alternative methods for measuring inorganic, organic, and total dissolved nitrogen in soil. *Soil Sci. Soc. Am. J.* 74:1018–1027. doi:10.2136/sssaj2009.0389
- Howarth, R., D. Swaney, G. Billen, J. Garnier, B. Hong, C. Humborg, P. Johnes, C.-M. Mörth, and R. Marino. 2012. Nitrogen fluxes from the landscape are controlled by net anthropogenic nitrogen inputs and by climate. *Front. Ecol. Environ* 10:37–43. doi:10.1890/100178
- Kim, D.G. 2008. Nitrous oxide and methane fluxes in riparian buffers and adjacent crop fields. Ph.D. diss., Iowa State University, Ames.
- Klemetsson, L., K. Von Arnold, P. Weslien, and P. Gundersen. 2005. Soil CN ratio as a scalar parameter to predict nitrous oxide emissions. *Glob. Change Biol.* 11:1142–1147. doi:10.1111/j.1365-2486.2005.00973.x
- Lin, C., R.N. Lerch, H.E. Garrett, D. Jordan, and M.F. George. 2007. Ability of forage grasses exposed to atrazine and isoxaflutole to reduce nutrient levels in soils and shallow groundwater. *Commun. Soil Sci. Plant Anal.* 38:1119–1136. doi:10.1080/00103620701327976
- Lowrance, R., R. Todd, J. Fail, O. Hendrickson, R. Leonard, and L. Asmussen. 1984. Riparian forests as nutrient filters in agricultural watersheds. *Bioscience* 34:374–377. doi:10.2307/1309729
- Matheson, F., M. Nguyen, A. Cooper, T. Burt, and D. Bull. 2002. Fate of ¹⁵N-nitrate in unplanted, planted and harvested riparian wetland soil microcosms. *Ecol. Eng.* 19:249–264. doi:10.1016/S0925-8574(02)00093-9
- Miller, W.P., and D.M. Miller. 1987. A micro-pipette method for soil mechanical analysis. *Commun. Soil Sci. Plant Anal.* 18:1–15. doi:10.1080/00103628709367799
- Mitchell, D.C., M.J. Castellano, J.E. Sawyer, and J. Pantoja. 2013. Cover crop effects on nitrous oxide emissions: Role of mineralizable carbon. *Soil Sci. Soc. Am. J.* 77:1765–1773. doi:10.2136/sssaj2013.02.0074
- Mitchell, D.C., X. Zhou, T.B. Parkin, M.J. Helmers, and M.J. Castellano. 2014. Comparing nitrate sink strength in perennial filter strips at toeslope of cropland watersheds. *J. Environ. Qual.* (in press).
- Mosier, A.R. 1980. Acetylene inhibition of ammonium oxidation in soil. *Soil Biol. Biochem.* 12:443–444. doi:10.1016/0038-0717(80)90023-1
- Mosier, A.R., and L. Klemetsson. 1994. Measuring denitrification in the field. In: R.W. Weaver, editor, *Methods of soil analysis. Part 2. SSSA Book Ser. 5. SSSA, Madison, WI.* p. 1047–1065.
- Myrold, D.D. 1990. Measuring denitrification in soils using ¹⁵N techniques. In: N.P. Revsbech and J. Sørensen, editors, *Denitrification in soil and sediment.* Plenum Press, New York. p. 181–198.
- Nestrud, L.M., and J.R. Worster. 1979. Soil survey of Jasper County, Iowa. USDA Soil Conservation Service and Iowa State University Cooperative Extension Service. U.S. Gov. Print. Office, Washington, DC.
- Parkin, T.B. 1987. Soil microsites as a source of denitrification variability. *Soil Sci. Soc. Am. J.* 51:1194–1199. doi:10.2136/sssaj1987.03615995005100050019x
- Parkin, T.B., H.F. Kaspar, A.J. Sexstone, and J.M. Tiedje. 1984. A gas-flow core method to measure field denitrification rates. *Soil Biol. Biochem.* 16:323–330. doi:10.1016/0038-0717(84)90026-9
- Parsons, W., M. Mitre, M. Keller, and W. Reiners. 1993. Nitrate limitation of N₂O production and denitrification from tropical pasture and rain forest soils. *Biogeochemistry* 22:179–193. doi:10.1007/BF00000646
- Pérez-Suárez, M., M.J. Castellano, R. Kolka, H. Asbjørnsen, and M. Helmers. 2014. Nitrogen and carbon dynamics in prairie vegetation strips across topographical gradients in mixed Central Iowa agroecosystems. *Agric. Ecosyst. Environ.* 188:1–11. doi:10.1016/j.agee.2014.01.023
- Peterjohn, W.T., and D.L. Correll. 1984. Nutrient dynamics in an agricultural watershed: Observations on the role of a riparian forest. *Ecology* 65:1466–1475. doi:10.2307/1939127
- Prather, M.J., R.D. Derwent, D. Ehhalt, P. Fraser, E. Sanhueza, and X. Zhou. 1995. Other trace gases and atmospheric chemistry. In: J.T. Houghton et al., editors, *Climate change 1994.* Cambridge Univ. Press, Cambridge, UK. p. 73–126.
- Prior, J.C. 1991. Landforms of Iowa. Univ. of Iowa Press, Iowa City.
- Reddy, K., and W. Patrick. 1984. Nitrogen transformations and loss in flooded soils and sediments. *Crit. Rev. Environ. Control* 13:273–309.
- Ryden, J.C., L.J. Lund, and D.D. Focht. 1979. Direct measurement of denitrification loss from soils: I. Laboratory evaluation of acetylene inhibition of nitrous oxide reduction. *Soil Sci. Soc. Am. J.* 43:104–110. doi:10.2136/sssaj1979.03615995004300010019x
- Ryden, J.C., J.H. Skinner, and D.J. Nixon. 1987. Soil core incubation system for the field measurement of denitrification in grassland soils. *Soil Biol. Biochem.* 19:753–757. doi:10.1016/0038-0717(87)90059-9
- Ryder, M.H., and A. Fares. 2008. Evaluating cover crops (sudex, sunn hemp, oats) for use as vegetative filters to control sediment and nutrient loading from agricultural runoff in a Hawaiian watershed. *J. Am. Water Resour. Assoc.* 44:640–653. doi:10.1111/j.1752-1688.2008.00189.x
- Schomberg, H.H., et al. 2009. Assessing indices for predicting potential nitrogen mineralization in soils under different management systems. *Soil Sci. Soc. Am. J.* 73:1575–1586. doi:10.2136/sssaj2008.0303
- Smith, M.S., and J.M. Tiedje. 1979. Phases of denitrification following oxygen depletion in soil. *Soil Biol. Biochem.* 11:261–267. doi:10.1016/0038-0717(79)90071-3
- Smith, P., et al. 2007. Policy and technological constraints to implementation of greenhouse gas mitigation options in agriculture. *Agric. Ecosyst. Environ.* 118:6–28. doi:10.1016/j.agee.2006.06.006
- Soil Survey Staff. 2003. *Keys to soil taxonomy.* 9th ed. USDA-NRCS, Washington, DC.
- Thorup-Kristensen, K., J. Magid, and L. Jensen. 2003. Catch crops and green manures as biological tools in nitrogen management in temperate zones. *Adv. Agron.* 79:227–302. doi:10.1016/S0065-2113(02)79005-6
- Zhou, X., M.J. Helmers, H. Asbjørnsen, R. Kolka, and M.D. Tomer. 2010. Perennial filter strips reduce nitrate levels in soil and shallow groundwater after grassland-to-cropland conversion. *J. Environ. Qual.* 39:2006–2015. doi:10.2134/jeq2010.0151
- Zhu, Q., and H.S. Lin. 2009. Simulation and validation of concentrated subsurface lateral flow paths in an agricultural landscape. *Hydrol. Earth Syst. Sci.* 13:1503–1518. doi:10.5194/hess-13-1503-2009
- Zhu, X., M. Burger, T.A. Doane, and W.R. Horwath. 2013. Ammonia oxidation pathways and nitrifier denitrification are significant sources of N₂O and NO under low oxygen availability. *Proc. Natl. Acad. Sci. USA* 110:6328–6333. doi:10.1073/pnas.1219993110